



Note

A note on binary matroid with no $M(K_{3,3})$ -minor

Xiangqian Zhou

Department of Mathematics, University of Mississippi, PO Box 1848, University, MS 38677, USA

Received 3 July 2003

Available online 25 April 2007

Abstract

We prove that if M is an internally 4-connected binary matroid with an $M(K_5)$ -minor and with no $M(K_{3,3})$ -minor, then either M has rank 4, or M is isomorphic to one of the following matroids: T_{12} , T_{12}/e , \widetilde{T}_{11} , \widetilde{T}_{12} , and \widetilde{T}_{13} .

© 2007 Elsevier Inc. All rights reserved.

Keywords: Binary matroids; Splitter Theorem; Blocking sequence; Internal 4-connectivity

1. Introduction

We assume the reader is familiar with matroid theory. Our notation and terminology will follow [5]. We begin by recalling Hall's Theorem [2].

Theorem 1.1 (Hall, 1943). *If G is a simple 3-connected graph, then G has no $K_{3,3}$ -minor if and only if G is planar or $G \cong K_5$.*

Theorem 1.1 strengthens Kuratowski's Theorem in the case of 3-connected graphs. It implies the only simple 3-connected graph with a K_5 -minor and with no $K_{3,3}$ -minor is the graph K_5 itself. We prove an analogue of Hall's Theorem for binary matroids in the internally 4-connected case.

Theorem 1.2. *Let M be an internally 4-connected binary matroid with an $M(K_5)$ -minor that has no $M(K_{3,3})$ -minor. Then either M has rank 4, or M is isomorphic to one of the following matroids: T_{12} , T_{12}/e , \widetilde{T}_{11} , \widetilde{T}_{12} , and \widetilde{T}_{13} .*

E-mail address: xzhou@olemiss.edu.

Note that there are precisely five 3-connected rank-4 binary matroids that have a proper $M(K_5)$ -minor. They are all internally 4-connected. The proof is a routine case checking. Therefore we have

Corollary 1.3. *There are precisely ten internally 4-connected binary matroids with a proper $M(K_5)$ -minor that have no $M(K_{3,3})$ -minor.*

To prove Theorem 1.2, we use Seymour's Splitter Theorem [7] and the blocking sequence techniques developed in [1] and [3]. A binary matroid will be represented by its fundamental graphs. The reader may refer to [6] or [8] for an introduction to these techniques.

The paper is constructed as follows: Section 2 contains the definitions of the matroids mentioned in Theorem 1.2 and in Section 3 we present the lemmas that are needed for the proof of Theorem 1.2.

2. Definitions

The matroid T_{12} was discovered by Kingan [4] as a splitter for the class of binary matroids with no $M(K_{3,3})$ - or $M^*(K_{3,3})$ -minor. The following matrix representation A_{12} of T_{12} was found by Oxley [4]:

$$A_{12} = \left[\begin{array}{c|cccccc} & 1 & 1 & 0 & 0 & 0 & 1 \\ & 1 & 0 & 0 & 0 & 1 & 1 \\ I_6 & 0 & 0 & 0 & 1 & 1 & 1 \\ & 0 & 0 & 1 & 1 & 1 & 0 \\ & 0 & 1 & 1 & 1 & 0 & 0 \\ & 1 & 1 & 1 & 0 & 0 & 0 \end{array} \right].$$

It is easy to see that T_{12} is self dual. T_{12} is 4-connected. Kingan has showed that T_{12} has a transitive automorphism group. Therefore, there is a unique single-element contraction (respectively deletion) of T_{12} , denoted by T_{12}/e (respectively $T_{12}\setminus e$). T_{12}/e and $T_{12}\setminus e$ are duals of each other and both are internally 4-connected.

The matroids \widetilde{T}_{11} , \widetilde{T}_{12} , and \widetilde{T}_{13} are represented by the binary matrices \widetilde{A}_{11} , \widetilde{A}_{12} , and \widetilde{A}_{13} , respectively:

$$\begin{aligned} \widetilde{A}_{11} &= \left[\begin{array}{c|cccccc} & 1 & 0 & 0 & 1 & 1 & 0 \\ & 1 & 1 & 0 & 0 & 0 & 1 \\ I_5 & 0 & 1 & 1 & 0 & 1 & 0 \\ & 0 & 0 & 1 & 1 & 0 & 1 \\ & 1 & 0 & 1 & 0 & 0 & 0 \end{array} \right], & \widetilde{A}_{12} &= \left[\begin{array}{c|cccccc} & 1 & 1 & 0 & 0 & 0 & 1 & 0 \\ & 1 & 0 & 0 & 0 & 1 & 1 & 1 \\ I_5 & 0 & 0 & 0 & 1 & 1 & 1 & 0 \\ & 0 & 0 & 1 & 1 & 1 & 0 & 1 \\ & 0 & 1 & 1 & 1 & 0 & 0 & 0 \end{array} \right], \\ \widetilde{A}_{13} &= \left[\begin{array}{c|cccccc} & 1 & 1 & 0 & 0 & 0 & 1 & 0 & 1 \\ & 1 & 0 & 0 & 0 & 1 & 1 & 1 & 0 \\ I_5 & 0 & 0 & 0 & 1 & 1 & 1 & 0 & 1 \\ & 0 & 0 & 1 & 1 & 1 & 0 & 1 & 1 \\ & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 \end{array} \right]. \end{aligned}$$

It is routine to check that the matroids \widetilde{T}_{11} , \widetilde{T}_{12} , and \widetilde{T}_{13} are internally 4-connected and have no $M(K_{3,3})$ -minor. Clearly $\widetilde{T}_{13} \setminus 13 = \widetilde{T}_{12}$.

Finally let Q_{10} be the binary matroid represented by the matrix A_{10} :

$$A_{10} = \left[\begin{array}{c|ccccc} & 1 & 1 & 0 & 0 & 0 \\ & 0 & 0 & 1 & 1 & 0 \\ I_5 & 1 & 0 & 1 & 0 & 1 \\ & 1 & 1 & 1 & 1 & 1 \\ & 0 & 1 & 0 & 1 & 1 \end{array} \right].$$

Note that Q_{10} is not internally 4-connected: $\{3, 4, 5, 10\}$ is a 4-element circuit and cocircuit. Q_{10} has an $M(K_{3,3})$ -minor but has no $M^*(K_{3,3})$ -minor.

3. Sketch of the proof

Let \mathcal{M} denote the class of binary matroid with no $M(K_{3,3})$ -minor. We rely heavily on the next lemma, which shows the matroid Q_{10}^* plays a similar role in \mathcal{M} to what the matroid R_{12} does in the class of regular matroids.

Lemma 3.1. *Let M be a binary matroid. If M has a Q_{10} -minor and M has no $M^*(K_{3,3})$ -minor; or equivalently, M has a Q_{10}^* -minor and has no $M(K_{3,3})$ -minor, then M is not internally 4-connected.*

It is easy to see that Theorem 1.2 follows from Lemmas 3.2–3.4.

Lemma 3.2. *Let M be a matroid in \mathcal{M} that has an $M(K_5)$ -minor. If M is 3-connected and has rank at least 5, then M has either a T_{12}/e - or a \widetilde{T}_{11} -minor.*

Lemma 3.3. *Let $M \in \mathcal{M}$. If M is 3-connected and has a proper T_{12}/e -minor, then M is isomorphic to T_{12} , \widetilde{T}_{12} , or \widetilde{T}_{13} .*

Lemma 3.4. *Let $M \in \mathcal{M}$ be internally 4-connected. If M has a \widetilde{T}_{11} -minor and has no T_{12}/e -minor, then M is isomorphic to \widetilde{T}_{11} .*

Lemmas 3.2–3.4 are proved by the same techniques as used in [6,8]. The proofs are straightforward but lengthy, thus omitted. A curious reader may look at [9] for details.

Acknowledgments

The author thanks Jim Geelen, Neil Robertson, Hongxun Qin and Geoff Whittle for many valuable discussions.

References

- [1] A. Bouchet, W.H. Cunningham, J.F. Geelen, Principal unimodular skew-symmetric matrices, *Combinatorica* 18 (1998) 461–486.
- [2] J.F. Geelen, private communication.
- [3] J.F. Geelen, A.M.H. Gerards, A. Kapoor, The excluded minors for $GF(4)$ -representable matroids, *J. Combin. Theory Ser. B* 79 (2001) 247–299.
- [4] S.R. Kingan, A generalization of a graph result of D.W. Hall, *Discrete Math.* 173 (1997) 129–135.
- [5] J.G. Oxley, *Matroid Theory*, Oxford Univ. Press, New York, 1992.

- [6] H. Qin, X. Zhou, The class of binary matroid with no $M(K_{3,3})$, $M^*(K_{3,3})$ -, $M(K_5)$ - or $M^*(K_5)$ -minor, *J. Combin. Theory Ser. B* 90 (2004) 173–184.
- [7] P.D. Seymour, Decomposition of regular matroids, *J. Combin. Theory Ser. B* 28 (1980) 305–359.
- [8] X. Zhou, On internally 4-connected non-regular binary matroids, *J. Combin. Theory Ser. B* 91 (2004) 327–343.
- [9] X. Zhou, Some excluded minor theorems for binary matroids, PhD dissertation, the Ohio State University, 2003.